

Lymphoma scanning

		Interim scan	End of chemo	Post RT/ RM	
Burkitt	No routine interim scanning in patients with clinical response	If response not assessable: -CT post 3x DA-EPOCH or 2x R-CODOX/M-IVAC -PET if predominantly extranodal	PET scan 6 weeks after last dose of chemo* (CT if prior CMR)		*disregard HD-MTX or intrathecal
DLBCL advanced	No routine interim scanning in patients with clinical response unless stipulated at MDT	If response not assessable: -CT post 3x R-CHOP -PET if bulky or predominantly extranodal	PET scan 6 weeks after last dose of chemo**	If RT: PET 3 months post RT (CT if prior CMR)	**3-4 weeks if HD MTX at EOT, or pre-planned consolidation RT not driven by PET response
DLBCL For CMT	Timely PET-3 RT to start within 3-6 weeks of last chemo	All patients: PET D15-18 of C3 R-CHOP	N/A#	PET 3 months post RT (CT if prior CMR)	# no repeat scan pre-RT if 4 th cycle R-CHOP given
PMBL R-CHOP	Routine iPET-3 RT to start within 3-6 weeks of last chemo	All patients: PET D10-13 of C3 R-CHOP-14 C4 to be continued without delay	No pre-RT scan if PET-3= CMR All others: PET scan 3 weeks after last dose of chemo	PET 3 months post RT (CT if prior CMR)	
PMBL DA-EPOCH	Routine iPET-3	All patients: PET D15-18 of C3 DA-EPOCH-R C4 to be continued without delay	PET scan 4 weeks after last dose of chemo (CT if prior CMR)	If RT (EOT PMR): PET 3 months post RT	
FL	No routine interim scanning in patients with clinical response	If response not assessable: -CT post 4x R-chemo -PET if predominantly extranodal	PET scan 6 weeks after last dose of chemo (no EOT scan if prior CMR)	PET scan at 12 months of RM (CT if prior CMR)	
MZL	No routine interim scanning in patients on R-chemo Routine CT after 4x R mono	R-chemo: if response not assessable: -CT post 4x R-chemo R mono: CT 4 weeks after 4 x weekly	CT scan 4 weeks after last dose of chemo/R mono (no EOT scan if prior CR)	CT at 12 months of RM	
MCL	No routine interim scanning in patients with clinical response unless auto-SCT planned	If response not assessable/planned SCT: -CT post 3x R-chemo -PET if predominantly extranodal	PET scan 6 weeks after last dose of chemo^ (CT if prior CMR)	PET scan at 12 months of RM (CT if prior CMR)	^3 weeks if SCT is planned